

Low Temperature Limit of Stability of Coherent Precession of Spin in the Superfluid $^3\text{He-B}$

E.V. Surovtsev, I.A. Fomin

P.L.Kapitza Institute for Physical Problems

Moscow Russia

July 13, 2009

Abstract

It is shown that instability of homogeneous precession is caused by combined effect of anisotropy of spin wave velocities and dipole interaction. In the principal order on the ratio of the Leggett frequency to the Larmor frequency the increments of growth of spin wave amplitudes are found. The magnitude of the maximum increment for all deviation angles of spin from its equilibrium orientation is calculated. The estimation is made of the minimum temperature down to which the precession is stable.

1. In a contrast to the superfluid A-phase, in the B-phase long wavelength perturbations do not destroy homogeneous precession of magnetization. Precession is stable with respect to such perturbation if the initial deviation angle of magnetization $\beta > \theta_0$, where $\theta_0 = \arccos(-1/4)$ and marginally stable if $\beta \leq \theta_0$. However, at low temperatures fast decay of homogeneous precession is observed in both cases. Such decay was first observed in experiments [1, 2] and is referred as *catastrophic relaxation*. Earlier we considered a process of parametric excitation of spin waves with finite wave vectors by the precession of magnetization as a possible origin of the decay [3]. In the theory of magnetics this effect is known as the *Suhl instability* [4]. The distinctive property of the instability in $^3\text{He-B}$ is that it occurs at the precession with large tipping angles $\beta \sim 100^\circ$ and also that excitation of different types of spin waves is possible. In Ref. [3] a scheme was proposed that takes into account both mentioned peculiarities and increments of instability for each of three types of spin waves were found. The dependence of the obtained

increments on magnetic field does not agree with that experimentally observed [5]. The magnitudes of the increments were also overvalued. The disagreement with the experiment was caused by our technical mistake. As a result of this mistake it turned out, that the instability of the precession can be induced by anisotropy of spin wave velocities in $^3\text{He-B}$ alone. The further analysis has shown that in order to provide coupling between spin waves and precession the dipole interaction has to be taken into account. It turned out also that considerable contribution to the instability comes from joint resonances when spin waves belonging to different branches of spectra are excited simultaneously.

In the present paper a revised version of the theory of parametric instability of homogeneous precession of spin in $^3\text{He-B}$ is presented. At the same time the foregoing disagreement with the experiment is resolved. A comparison is also made with the results of Ref.[6, 7], where contribution of the boundaries to the development of parametric instability is considered.

2. Following the procedure of Ref.[3] let us parameterize orientation of the order parameter of $^3\text{He-B}$, which is a rotation matrix $R_{\xi i}$, by the Euler angles α, β, γ (z -axis is oriented opposite to the direction of the d.c. magnetic field $\mathbf{H}_0$). Actually, it is more convenient to use the sum $\Phi = \alpha + \gamma$ instead of the angle γ . Canonically conjugated momenta to these coordinates are correspondingly the following combinations of spin projections $P = S_z - S_\zeta, S_\beta, S_\zeta$, where S_z — is projection of spin onto z -axis, S_ζ — its projection onto $\zeta = \hat{R}\hat{z}$ and S_β — is projection on the line of nodes (see for example [8]). Equations of motion are hamiltonian with respect to the stated pairs of variables with the Hamiltonian

$$H = \frac{1}{1 + \cos \beta} \left\{ S_\zeta^2 + P S_\zeta + \frac{P^2}{2(1 - \cos \beta)} \right\} + \frac{1}{2} S_\beta^2 + F_\nabla - \omega_L (P + S_\zeta) + U_D(\alpha, \beta, \Phi). \quad (1)$$

Here ω_L — is the Larmor frequency, corresponding to the d.c. magnetic field, F_∇ — the gradient energy, $U_D(\alpha, \beta, \Phi)$ — the dipole energy, which is of the order of the squared Leggett frequency in the $^3\text{He-B}$: Ω_B^2 . We choose the units of measurements so that the magnetic susceptibility of $^3\text{He-B}$ — χ , and the gyromagnetic ratio for nuclei of ^3He — g are equal to unity. In this units spin has a dimensionality of frequency and energy — of a squared frequency correspondingly. In the standard setting of NMR experiments in the regions distant from the walls of the cell spin precesses in the so called *Leggett configuration*, when the "orbital vector" $l_i = -R_{\xi i} s_\xi$ [6] is parallel to the magnetic field. In this case $U_D(\alpha, \beta, \Phi)$ does not depend on the angle α and precession is described by the stationary solution of equations of spin

dynamics which do not contain oscillating terms:

$$\begin{aligned} \alpha &= -\omega_p t + \alpha_0, \quad \gamma = \omega_p t + \Phi^{(0)} - \alpha_0, \\ P^{(0)} &= \omega_p (\cos \beta - 1), \quad S_\beta^{(0)} = 0, \quad S_z^{(0)} = \omega_p \cos \beta, \end{aligned} \quad (2)$$

where ω_p – the frequency of precession. If $\beta > \theta_0$ then $\Phi^{(0)} = 0$, and if $\beta < \theta_0$ –

$$\cos \Phi^{(0)} = (\frac{1}{2} - \cos \beta^{(0)}) / (1 + \cos \beta^{(0)}). \quad (3)$$

Explicit time dependence of the stationary solution (2) can be excluded if one transfers to the variable $\psi = \alpha + \omega_p t$ and uses new Hamiltonian $\tilde{H} = H + \omega_p P$, then $\frac{\partial \psi}{\partial t} = 0$. To find the spectra of excitation against the background of precession we linearize the equations of motion on small deviations from the stationary solution (2, 3): $\delta\psi(\mathbf{r}, t) = \psi - \psi^{(0)}$, etc. For the sake of convenience the following combinations of the mentioned deviations are used:

$$\begin{aligned} \nu &= \sin \beta \delta\psi, \quad \vartheta = \frac{\delta P + (1 - \cos \beta) \delta S_\zeta}{\omega_L \sin \beta}, \\ \varepsilon &= \delta \Phi - (1 - \cos \beta) \delta\psi, \quad \sigma = \delta S_\zeta / \omega_L, \\ \zeta &= \delta S_\beta / \omega_L, \quad \eta = -\delta\beta. \end{aligned} \quad (4)$$

Expression for the gradient energy of ${}^3\text{He} - B$ contains two coefficients that can be written as velocities of two types of spin waves $c_\parallel$ and $c_\perp$. In what follows units of length and time are chosen so that $\omega_L = 1$ and $c_\parallel = 1$. Without loss of generality one can assume that variables change only in y and z directions, then the time-independent part of the gradient energy has the form:

$$F_{\nabla st} = \frac{1}{2} [(1 - \mu)(\nu_{,y}^2 + \eta_{,y}^2) + \nu_{,z}^2 + \eta_{,z}^2 + \varepsilon_{,y}^2 + (1 - 2\mu)\varepsilon_{,z}^2], \quad (5)$$

where $\mu = 1 - c_\perp^2 / c_\parallel^2$ – is the anisotropy of spin wave velocities. Parameter μ will be considered as a small one, in fact $\mu \approx 1/4$ [9]. Furthermore, F_∇ also contains term oscillating with the frequency of precession:

$$F_{\nabla osc1} = -\mu [(\eta_{,y}\varepsilon_{,z} + \eta_{,z}\varepsilon_{,y}) \cos \omega_p t + (\nu_{,y}\varepsilon_{,z} + \nu_{,z}\varepsilon_{,y}) \sin \omega_p t] \quad (6)$$

and with the doubled frequency of precession

$$F_{\nabla osc2} = -\frac{\mu}{2} [(\eta_{,y}^2 - \nu_{,y}^2) \cos 2\omega_p t + 2\nu_{,y}\eta_{,y} \sin 2\omega_p t]. \quad (7)$$

In zero order approximation on small parameters μ and $(\Omega_B / \omega_L)^2$ the equations for deviations have hamiltonian form with the Hamiltonian:

$$h = \frac{1}{2} [(\vartheta + \eta)^2 + \zeta^2 + \sigma^2 + (\nabla \nu)^2 + (\nabla \eta)^2 + (\nabla \varepsilon)^2] \quad (8)$$

and with respect to the pairs of canonically conjugated variables (ε, σ) ; (ν, ϑ) ; (ζ, η) . In each pair the first variable is coordinate and the second is momentum. Equations of motion for the pair (ε, σ) have the form:

$$\frac{\partial \varepsilon}{\partial t} = \sigma, \quad \frac{\partial \sigma}{\partial t} = \Delta \varepsilon. \quad (9)$$

It is convenient to rewrite them in a vectorial form:

$$\frac{d\mathbf{X}_1}{dt} = \hat{M}_1 \mathbf{X}_1. \quad (10)$$

Solutions of the system (9) have the form of plane waves $\mathbf{e}_1^\pm \exp[i(\mathbf{k}\mathbf{r} \mp \omega_1 t)]$ with the dispersion law

$$\omega_1 = k. \quad (11)$$

Here $\mathbf{e}_1^\pm$ are right eigenvectors of matrix $\hat{M}_1$ corresponding to eigenvalues $\mp i\omega_1$:

$$\mathbf{e}_1^\pm = \begin{pmatrix} 1 \\ \mp i\omega_1 \end{pmatrix}. \quad (12)$$

One needs left eigenvectors $\mathbf{f}_1^\pm$ of the same matrix to be able to make projections. They can be normalized so that the following conditions are met:

$$(\mathbf{f}_1^+, \mathbf{e}_1^+) = 1; \quad (\mathbf{f}_1^+, \mathbf{e}_1^-) = 0. \quad (13)$$

Here the scalar product is defined as

$$(\mathbf{a}, \mathbf{b}) = \sum_{n=1,2} a_n^* b_n. \quad (14)$$

As a result

$$\mathbf{f}_1^\pm = \frac{1}{2\omega_1} (\omega_1; \mp i). \quad (15)$$

In a similar way the following dispersion laws for two transverse modes are obtained:

$$\omega_2 = \sqrt{\frac{1}{4} + k^2} - \frac{1}{2}, \quad (16)$$

$$\omega_3 = \sqrt{\frac{1}{4} + k^2} + \frac{1}{2}. \quad (17)$$

At $k \rightarrow 0$ $\omega_2 \sim k^2$, i.e. it is a gapless mode arising from the degeneracy of precession with respect to α_0 . Another mode has a gap $\omega_3 = \omega_L$ at

$k \rightarrow 0$, it passes into nutations. In the coordinates $\psi, \vartheta, \eta, \zeta$ the following right eigenvectors correspond to the frequencies $\pm\omega_2$:

$$\mathbf{e}_2^\pm = \begin{pmatrix} 1 \\ \mp i\omega_3 \\ \pm i \\ \omega_2 \end{pmatrix}, \quad (18)$$

and to the frequencies $\pm\omega_3$:

$$\mathbf{e}_3^\pm = \begin{pmatrix} 1 \\ \mp i\omega_2 \\ \mp i \\ -\omega_3 \end{pmatrix}. \quad (19)$$

Left eigenvectors are correspondingly

$$\mathbf{f}_2^\pm = \frac{1}{2\omega_{23}} (\omega_3; \mp i; \pm i\omega_2; 1), \quad (20)$$

and

$$\mathbf{f}_3^\pm = -\frac{1}{2\omega_{23}} (-\omega_2; \pm i; \pm i\omega_3; 1), \quad (21)$$

where $\omega_{23} = \omega_2 + \omega_3$.

3. Time-dependent corrections to the Hamiltonian (8) can provide creation and mutual transformation of excitations. At $\mathbf{l} \parallel \mathbf{H}_0$ the dipole energy does not contain time-dependent terms and the gradient energy in the first approximation on μ contains oscillating terms (6),(7). Taking the oscillating terms into account one can write equations of motion for deviations $\varepsilon, \sigma, \nu, \vartheta, \zeta, \eta$ combined in a six-component vector-column in a form:

$$\frac{d\mathbf{X}}{dt} = (\hat{M}_0 + \hat{V}(t)) \mathbf{X}, \quad (22)$$

where all time-dependent terms are collected in $\hat{V}(t)$. The sum of (6) and (7) yields $\hat{V}(t) = \sum_{n=1,2} [\hat{W}_n \exp(-in\omega_p t) + \hat{W}_n^* \exp(in\omega_p t)]$. Following procedure of time-dependent perturbation theory let us seek for a solution of the system (22) in a form of expansion in eigenvectors of matrix $\hat{M}_0$

$$\mathbf{X}(\mathbf{r}, t) = \sum_{j, \mathbf{k}} \{a_{j\mathbf{k}}^+(t) \mathbf{e}_{j\mathbf{k}}^+ \exp(i\mathbf{k}\mathbf{r} - i\omega_j t) + a_{j\mathbf{k}}^-(t) \mathbf{e}_{j\mathbf{k}}^- \exp(i\mathbf{k}\mathbf{r} + i\omega_j t)\} \quad (23)$$

Eigenvectors $\mathbf{e}_{j\mathbf{k}}^\pm$ are the mentioned above $\mathbf{e}_1^\pm, \mathbf{e}_2^\pm, \mathbf{e}_3^\pm$ supplemented with zeroes up to six component. Substitution of (23) into (22) and separation of

equations by $\mathbf{k}$ yields:

$$\sum_j \{ \dot{a}_{j\mathbf{k}}^+ \mathbf{e}_{j\mathbf{k}}^+ \exp(-i\omega_j t) + \dot{a}_{j\mathbf{k}}^- \mathbf{e}_{j\mathbf{k}}^- \exp(i\omega_j t) \} = \sum_{j,n} (\hat{W}_n \exp(-in\omega_p t) + \hat{W}_n^* \exp(in\omega_p t)) \{ a_{j\mathbf{k}}^+ \mathbf{e}_{j\mathbf{k}}^+ \exp(-i\omega_j t) + a_{j\mathbf{k}}^- \mathbf{e}_{j\mathbf{k}}^- \exp(i\omega_j t) \} \quad (24)$$

Multiplying two sides of equation (24) by $\mathbf{f}_l^+ \exp(i\omega_l t)$ and omitting common index $\mathbf{k}$ one has:

$$\dot{a}_l^+ = \sum_{j,n} \{ a_j^- (\mathbf{f}_l^+, \hat{W}_n \mathbf{e}_j^-) \exp[i(\omega_l + \omega_j - n\omega_p)t] + a_j^+ (\mathbf{f}_l^+, \hat{W}_n^* \mathbf{e}_j^+) \exp[i(\omega_l - \omega_j + n\omega_p)t] \}. \quad (25)$$

Application of the procedure assumes that coefficients $a_{j\mathbf{k}}^\pm$ are weakly varying at time $\sim 1/\omega_e$. Averaging of equation (25) shows that non-trivial correlations between different $a_{j\mathbf{k}}^\pm$ arise only nearby the resonances $\omega_j(\mathbf{k}) - \omega_l(\mathbf{k}) = n\omega_p$ and $\omega_l(\mathbf{k}) + \omega_j(-\mathbf{k}) = n\omega_p$. The second resonance corresponds to creation of quasiparticles from "vacuum". In accordance with equality $\omega_j(\mathbf{k}) = \omega_j(-\mathbf{k})$ the sign of one of the momenta is changed. In what follows it is assumed that the resonance condition is fulfilled exactly then the obtained increment of instability is a maximum one. If at given $\mathbf{k}$ the resonance condition is fulfilled only for two states l and j , then

$$\dot{a}_l^+ = (\mathbf{f}_l^+, \hat{W} \mathbf{e}_j^-) a_j^-. \quad (26)$$

The same argument for a_j^- yields:

$$\dot{a}_j^- = (\mathbf{f}_j^-, \hat{W}^* \mathbf{e}_l^+) a_l^+. \quad (27)$$

The system of equations (26),(27) has solutions $\sim \exp(\pm \lambda t)$, where λ is determined by $\lambda^2 = (\mathbf{f}_j^+, \hat{W} \mathbf{e}_l^-) (\mathbf{f}_l^+, \hat{W} \mathbf{e}_j^-)^*$. Thus the problem of determination of the increment of instability reduces to calculation of the elements of matrix $(\mathbf{f}_j^+, \hat{W} \mathbf{e}_l^-)$ between the states satisfying the resonance conditions. Particularly, if $l = j$, then $2\omega_l(\mathbf{k}) = n\omega_p$ and $\lambda^2 = |(\mathbf{f}_j^+, \hat{W} \mathbf{e}_l^-)|^2$. This is the simplest case of *parametric resonance* [10]. Explicit expressions for $\hat{W}_1$ and $\hat{W}_2$ are found using equalities (6) and (7). There are four non-zero elements of matrix $\hat{W}_1$: $(\hat{W}_1)_{\sigma\nu} = (\hat{W}_1)_{\vartheta\varepsilon} = i\mu k_y k_z$ and $(\hat{W}_1)_{\sigma\eta} = (\hat{W}_1)_{\zeta\varepsilon} = \mu k_y k_z$, and there are another four finite elements of matrix $\hat{W}_2$: $(\hat{W}_2)_{\vartheta\nu} = -(\hat{W}_2)_{\zeta\eta} = \mu k_y^2/2$ and $(\hat{W}_2)_{\vartheta\eta} = (\hat{W}_2)_{\zeta\nu} = i\mu k_y^2/2$. Here k_y, k_z are the components of the wave vector satisfying the resonance conditions. Direct verification shows that the matrix element $(\mathbf{f}_j^+, \hat{W}_1 \mathbf{e}_l^-)$ is not equal to

zero only if $j = 1, l = 3$. The resonance condition $\omega_1(\mathbf{k}) + \omega_3(-\mathbf{k}) = \omega_L$ is fulfilled only at $k = 0$, but at this value of k the matrix element turns into zero due to the factor $k_y k_z$. For the matrix $(\mathbf{f}_j^+, \hat{W}_2 \mathbf{e}_l^-)$ non-zero element corresponds to $j = l = 3$. In this case the resonance condition $\omega_3(\mathbf{k}) + \omega_3(-\mathbf{k}) = 2\omega_L$ is also fulfilled only at $k = 0$ and the corresponding matrix element is equal to zero due to the factor k_y^2 . Thus if the dipole energy is neglected then the anisotropy of spin wave velocities does not provide the coupling between precession and spin waves.

4. In order to take the dipole energy into account it is need to add to the RHS of equation (22) the matrix of dipole torque $\hat{N}_D$ and to find new energies of excitations and the corresponding eigenvectors. The added terms in the equation (22) are small in comparison with the elements of the matrix $\hat{M}_0$ as $(\Omega_B/\omega_L)^2$, which is in typical experimental conditions of the order of $\sim 10^{-1} - 10^{-2}$. In the first order approximation on the mentioned parameter the elements of matrix $(\mathbf{f}_j^+, \hat{W}_n \mathbf{e}_l^-)$ are given by the expressions:

$$\begin{aligned} (\mathbf{f}_j^+, \hat{W}_n \mathbf{e}_l^-) = & i \sum_{m \neq j+} \frac{1}{\omega_{j+} - \omega_m} (\mathbf{f}_j^+, \hat{N}_D \mathbf{e}_m) (\mathbf{f}_m, \hat{W}_n \mathbf{e}_l^-) + \\ & i \sum_{m \neq l-} \frac{1}{\omega_{l-} - \omega_m} (\mathbf{f}_j^+, \hat{W}_n \mathbf{e}_m) (\mathbf{f}_m, \hat{N}_D \mathbf{e}_l^-). \end{aligned} \quad (28)$$

Non-zero elements are obtained at the following resonances:

$$\omega_1(k) + \omega_1(-k) = \omega_p, \quad k = 1/2, \quad \beta \leq \theta_0, \quad (29)$$

$$\omega_1(k) + \omega_2(-k) = \omega_p, \quad k = 2/3, \quad (30)$$

$$\omega_1(k) + \omega_3(-k) = 2\omega_p, \quad k = 2/3, \quad (31)$$

$$\omega_2(k) + \omega_3(-k) = 2\omega_p, \quad k = \sqrt{3}/2. \quad (32)$$

The corresponding increments are of the order of $\mu\Omega_B^2/\omega_L$. At $k=1/2$

$$\lambda_{(11)} = \begin{cases} \mu \sin(2\delta) \frac{\sqrt{(1 - \cos(\beta))(1 + 4 \cos(\beta))} \Omega_B^2}{5 \omega_L}, & \beta \leq \theta_0, \\ 0 & \beta > \theta_0, \end{cases} \quad (33)$$

where δ is the angle between the direction of the wave vector $\mathbf{k}$ and the direction of the magnetic field. At $k=2/3$ there are two resonances and one needs to consider the system of equations for thee amplitudes in order to find the increment. In this case the increment is given by the expression

$$\lambda_{(12,13)} = \sqrt{\lambda_{(12)}^2 + \lambda_{(13)}^2}, \quad (34)$$




Figure 1:
Dependence of the coefficient a in formule (38) on tipping angle β

where

$$\lambda_{(12)} = \begin{cases} \mu \sin(2\delta) \frac{\sqrt{10}(1 - \cos(\beta)) \Omega_B^2}{25 \omega_L}, & \beta \leq \theta_0, \\ \mu \sin(2\delta) \frac{|4 \cos^2(\beta) + 31 \cos(\beta) + 15| \Omega_B^2}{15\sqrt{10} \omega_L}, & \beta > \theta_0, \end{cases} \quad (35)$$

$$\lambda_{(13)} = \begin{cases} \mu \sin^2(\delta) \frac{\sqrt{10} \sqrt{(1 - \cos(\beta))(1 + 4 \cos(\beta))} \Omega_B^2}{25 \omega_L}, & \beta \leq \theta_0, \\ -\mu \sin^2(\delta) \frac{\sqrt{10} \sin(\beta)(1 + 4 \cos(\beta)) \Omega_B^2}{225 \omega_L}, & \beta > \theta_0, \end{cases} \quad (36)$$

The increment for the resonance at $k = \sqrt{3}/2$ is

$$\lambda_{(23)} = \begin{cases} \mu \sin^2(\delta) \frac{3(1 - \cos(\beta)) \Omega_B^2}{20 \omega_L}, & \beta \leq \theta_0, \\ \mu \sin^2(\delta) \frac{|4 \cos^2(\beta) + 31 \cos(\beta) + 15| \Omega_B^2}{40 \omega_L}, & \beta > \theta_0. \end{cases} \quad (37)$$

The maximum increment for each value of β can be written as

$$\lambda_{max}(\beta) = a(\beta) \mu \frac{\Omega_B^2}{\omega_L}, \quad (38)$$

where the dependence of the coefficient a on tipping angle is shown at Fig.1.

At finite temperatures the damping of spin waves has to be taken into account. The instability sets up if the increment of growth of waves, which satisfy the resonance condition, exceeds the decrement of damping. As before

[3] for estimation of the temperature of catastrophic relaxation T_{cat} it will be assumed here that the principal mechanism of dissipation is spin diffusion. The minimum temperature, down to which the precession is stable, is found from the equation:

$$\frac{D(T)k^2}{2} = \lambda_{max}, \quad (39)$$

where $D(T)$ is the coefficient of spin diffusion. At temperatures in question $T \leq 0.4T_c$, the increment weakly depends on temperature and its value can be taken at $T = 0$. The LHS of (39) strongly depends on temperature owing to spin diffusion which behaves as $D(T) \sim \sqrt{T/\Delta} \exp(-\Delta/T)$ at $T \rightarrow 0$.

The obtained increments originate from the coupling between precession and spin waves in bulk helium. In Ref. [6, 7] the increment arising from enhancement of the coupling in regions adjacent to the walls was found. Because of boundary conditions precession on the walls goes on in the configuration different from the Leggett one. In this case oscillating terms in the dipole energy appear without taking the anisotropy of spin waves velocities into account. The local coupling in regions adjacent to the walls is on the order of $1/\mu$ greater than that obtained for the bulk helium. In Ref.[6, 7] the result of calculation of increment of growth and T_{cat} for the angle $\beta = 90^\circ$ is represented. The bulk contribution to the increment for the same conditions estimated with the use of the above formulae gives approximately the same value as the surface contribution. It is impossible to separate surface and bulk contributions to the increment by their dependence on magnetic field because both contributions are proportional to Ω_B^2/ω_L . However, it has to be mentioned, that the surface contribution depends on ratio of volume adjacent to the surfaces to the total volume of helium. For angles $\beta < 104^\circ$ there is no characteristic length whereon penetrates the perturbation effect of the walls on the precession. Thus the regions adjacent to the walls occupies considerable part of the volume. The most part of the data about catastrophic relaxation is obtained in experiments with the homogeneously precessing domain. In this case magnetization precesses with angles slightly above $\theta_0 \approx 104^\circ$ and the frequency of precession is shifted from the Larmor frequency. Then the effect of the walls is limited by the "coherence length" $\xi = c_\perp/\sqrt{\omega_p(\omega_p - \omega_L)}$ [11, 12]. For the typical experimental conditions in the most part of the precessing domain $\xi \sim 10^{-2}$ cm and the regions adjacent to the walls occupies only a small part of the total volume. The increment is determined by the bulk resonance $\omega_2(k) + \omega_3(-k) = 2\omega_p$. The available experimental data for diffusion coefficient [13] pertains to the temperatures $T > 0.4T_c$. In the experiments [1] at pressure $P \simeq 20$ bar and magnetic field $H \simeq 142$ Oe $T_{cat} \simeq 0.42T_c$. After substitution of the diffusion coefficient

at this temperature $D = 0.04 \text{ cm}^2/c$ and the values of other parameters $\Omega = 2\pi \cdot 250 \text{ kHz}$, $c_{\parallel} = 1.6 \cdot 10^3 \text{ cm/c}$, $\mu = 1/4$ into the formula (39) we obtain for the LHS of equation (39) the value $5 \cdot 10^4 1/c$, and for the RHS $-4 \cdot 10^4 1/c$. Taking into account that parameters μ and Ω_B^2/ω_L^2 are not too small the obtained agreement can be regarded as a satisfactory for this example.

5. In conclusion the Suhl instability limits from below the interval of temperatures where coherent precession in $^3\text{He}-B$ can exist. The obtained here low temperature limit of stability of precession is caused by the interaction of precession with spin waves in the bulk helium.

Lowering of the limit temperature can be achieved by using higher magnetic fields as it was demonstrated in experiments [5]. Such tendency agrees with formula (39). On the one hand increasing of magnetic field decreases the increment of instability and on the other hand it increases spin waves damping. However, due to the exponential dependence of the coefficient of diffusion on temperature the effect of magnetic field on T_{cat} becomes weaker when temperature decreases. Detailed comparison with the results of [5] will be done in a full-length publication.

We thank to V.V. Dmitriev for useful discussion. This research was supported by RFBR and Ministry of Education and Science of Russian Federation.

References

- [1] Yu.M. Bunkov, V.V. Dmitriev, Yu.M. Mukharsky et al., *Europhysics Lett.* **8**, 645 (1989).
- [2] Yu.M. Bunkov, V.V. Dmitriev, J. Nyeki et al., *Physica B* **165**, 675 (1990).
- [3] E.V.Surovtsev, I.A. Fomin, *Pis'ma ZheTF* **83**, 479 (2006).
- [4] H. Suhl, *J. Phys. Chem. Solids*, **1**, 209 (1957).
- [5] D.A. Geller and D.M. Lee, *Phys. Rev. Lett.* **85**, 1032 (2000).
- [6] Yu.M. Bunkov, V.S. Lvov, G.E. Volovik, *Pis'ma v ZhETF* **83**, 624 (2006)
- [7] Yu.M. Bunkov, V.S. Lvov, G.E. Volovik, *Pis'ma v ZhETF* **84**, 349 (2006)
- [8] I. A. Fomin, *Zh. Exp. Teor. Fiz.* **84**, 2109 (1983) [*Sov. Phys. JETP* **57**, 1227 (1983)].

- [9] Yu.M. Bunkov, V.V. Dmitriev, Yu.M. Mukharsky, Pis'ma ZheTF **43**, 131 (1986)
- [10] L. D. Landau and E. M. Lifshitz, *Course of Theoretical Physics*, Vol. 1: *Mechanics*, 4th ed. (Nauka, Moscow, 1988, Pergamon, Oxford, 1989)
- [11] I. A. Fomin, Zh. Exp. Teor. Fiz. **94**, 112 (1988).
- [12] Yu.M. Bunkov, O.D. Timofeevskaya, G.E. Volovik, Phys. Rev. Lett. **73**, 1817 (1994).
- [13] Yu.M. Bunkov, V.V. Dmitriev, A.V. Markelov et al., Phys. Rev. Lett. **65**, 867 (1990).